



POST-SURGERY INSTRUCTIONS: BREAST LIFT with SALINE IMPLANTS

Once your surgery is completed, you must follow all the instructions given to you in order to heal properly and have a good outcome.

The following instructions are your obligation. Use this as a checklist of progress as you heal. Included are normal post-surgical experiences and key health considerations that may be a cause of concern.

TYPICAL POST-OPERATIVE SYMPTOMS

Typical symptoms of breast surgery and signs to watch for following breast lift surgery with saline filled implants include the following:

Tightness in the chest region and stiffness: Tingling, burning, or intermittent shooting pain. These are normal experiences as the skin, muscles, and tissue stretch to accommodate your implants, and as sensory nerves heal. Pain medication and muscle relaxants will help you cope with any discomfort. If you have drains, you may experience additional localized discomfort. **Consistent sharp pain should be reported to our office immediately.**

Hypersensitivity of nipples or lack of sensitivity: This is normal and will gradually resolve over time. You may also experience a small amount of fluid or milk seeping through the nipples. **If this becomes painful or excessive notify our office immediately.**

A feeling of heaviness: It is normal for your chest to feel heavy as you adjust to your implants. This will subside within 2-4 weeks following surgery.

Shiny skin or any itchy feeling: Swelling can cause the breasts skin to appear shiny. As the healing process advances, you may also find a mild to severe itchy feeling of the breasts. An antihistamine like Benadryl can help to alleviate severe, constant itchiness. **If the skin becomes red and hot to the touch, contact our office immediately.**

Asymmetry, the breasts look different, or heal differently. Breasts may look or feel quite different from one another in the days following surgery. This is normal. Although no two breasts are perfectly symmetrical in nature or following surgery, breast massage and time will produce breasts that are similar in shape, position, and size.

A sloshing sound or sensation: This is not the result of your saline implant filler, but rather of air that is trapped in the implant pocket and fluid that may naturally accumulate. This is perfectly normal and will resolve within 2-4 weeks.

CALL THE OFFICE IMMEDIATELY IF YOU EXPERIENCE ANY OF THE FOLLOWING:

- **A high fever, (over 101°) severe nausea and vomiting, continued dizziness or incoherent behavior, such as hallucinations.**
- **Any pain that cannot be controlled by your pain medication.**
- **Bright red skin that is hot to the touch.**
- **Excessive bleeding or fluid seeping through the incisions.**
- **A severely misshapen breast or bruising that is localized to one breast or region of the chest.**

DAY OF SURGERY INSTRUCTIONS:

- Walk every 2 hours for 10min, while awake
- Resume daily activities as tolerated
- Take medications as directed
- You may resume all medications immediately after surgery
- Sleep at 45 degree angle (2 pillows) for 1-2 weeks
- Keep incision clean and dry.
- Do not remove steri-strips (little white or brown tape strips)
- You may shower 24-48 hours after surgery.
- Do Not Drive while taking a narcotic of for 72hr post surgery.
- You may switch to Ibuprofen 24 hours after procedure
- Wear compression garment as directed
- Do not lift, pull, or push anything heavier than 5lbs for 2 weeks post surgery
- No exercising for 14 days after surgery
- Schedule follow up appointment 3-5 days post surgery

To contact Dr. Few after hours, please call the answering service at 312-202-0882 , and press "0"